

## $^7\text{Li}$ Ion Diffusion in Isotope-diluted Glassy $\text{Li}_2\text{Si}_3\text{O}_7$ – The Generation of pure Spin-3/2 Spin-alignment NMR Echoes

Dominik Wohlmuth<sup>1</sup>, Viktor Epp<sup>1</sup>, Ute Bauer<sup>1</sup>, Anna-Maria Welsch<sup>2</sup>, Harald Behrens<sup>2</sup>,  
Martin Wilkening<sup>1</sup>

<sup>1</sup> Institute for Chemistry and Technology of Materials, Christian Doppler Laboratory for Lithium Batteries, Graz University of Technology, Stremayrgasse 9, 8010 Graz, Austria; <sup>2</sup> Institut für Mineralogie, Leibniz Universität Hannover, Callinstr. 3, 30167 Hannover, Germany

E-Mail: wilkening@tugraz.at

Solid-state diffusion plays one of the most important roles in materials science. In particular, the precise measurement of ion dynamics in materials with structural disorder is of great interest. Spin-alignment echo (SAE) nuclear magnetic resonance (NMR), being comparable to exchange spectroscopy, turned out to be a powerful method to probe (ultra-)slow Li dynamics even in amorphous materials [1–3]. However,  $^7\text{Li}$  Jeener-Broekaert echoes can be influenced by the simultaneous generation of dipolar with quadrupolar order. In many cases, the first can be suppressed by choosing proper evolution times  $t_p$  of less than 20  $\mu\text{s}$  [4, 5].

Here, glassy  $\text{Li}_2\text{Si}_3\text{O}_7$  served as a suitable model system to study the positive influence of isotope dilution on  $^7\text{Li}$  SAE NMR, *i.e.*, the reduction of homonuclear dipole-dipole interactions through spatial separation of the spin-3/2 probe nuclei. Two samples, one with 100%  $^7\text{Li}$  and the other one with 5%  $^7\text{Li}$  (95%  $^6\text{Li}$ ), were investigated by  $^7\text{Li}$  NMR line-shape analysis, spin-lattice relaxation NMR as well as mixing-time and preparation-time dependent  $^7\text{Li}$  SAE NMR using a 32-fold phase cycle. Jeener-Broekaert echoes and their Fourier transforms show that at sufficiently short  $t_p$  the interfering dipolar interactions can be completely suppressed in that sample for which the proportion of  $^7\text{Li}$  was greatly reduced by substitution with  $^6\text{Li}$ . The so-obtained diffusion parameters are compared with results deduced from broadband conductivity spectroscopy.

**Acknowledgement.** We thank our colleagues at the University of Hannover and the TU Graz for valuable discussions. Financial support by the Deutsche Forschungsgemeinschaft (WI3600/2-2 and 4-1) as well as by the Austrian Federal Ministry of Science, Research and Economy, and the Austrian National Foundation for Research, Technology and Development is greatly appreciated.

- [1] Studying Li Dynamics in a Gas-Phase Synthesized Amorphous Oxide by NMR and Impedance Spectroscopy, V. Epp, C. Brüning, M. Binnewies, P. Heitjans, M. Wilkening, *Z. Phys. Chem.* 226 (2012) 513.
- [2] Atomic-scale Measurement of Ultraslow Li Motions in Glassy  $\text{LiAlSi}_2\text{O}_6$  by two-time  $^6\text{Li}$  Spin-alignment Echo NMR Correlation Spectroscopy, M. Wilkening, A. Kuhn, P. Heitjans, *Phys. Rev. B* 78 (2008) 054303.
- [3] From Micro to Macro: Access to Long-Range  $\text{Li}^+$  Diffusion Parameters in Solids via Microscopic  $^{6,7}\text{Li}$  Spin-Alignment Echo NMR Spectroscopy, M. Wilkening, P. Heitjans, *Chem. Phys. Chem.* 13 (2012) 53.
- [4] Alignment Echo of Spin-3/2  $^9\text{Be}$  Nuclei: Detection of Ultraslow Motion, X.-P. Tang and Y. Wu, *J. Magn. Res.* 133 (1998) 155.
- [5] Ion Dynamics in Solid Electrolytes: NMR Reveals the Elementary Steps of  $\text{Li}^+$  Hopping in the Garnet  $\text{Li}_{6.5}\text{La}_3\text{Zr}_{1.75}\text{Mo}_{0.25}\text{O}_{12}$ , P. Bottke, D. Rettenwander, W. Schmidt, G. Amthauer, M. Wilkening, *Chem. Mater.* 27 (2015) 6571.

